



Case Files In Glaucoma

Jessica Steen, O.D.,F.A.A.O.

Eric Schmidt, O.D.,F.A.A.O.



JESSICA STEEN OD FINANCIAL DISCLOSURES

- Speakers Bureau-Carl Zeiss Meditec, Bausch and Lomb, Viatris, Thea Pharma, Alcon, Allergan, Astellas, Dompé
- Consultant-Bausch and Lomb, Balance Ophthalmics, Carl Zeiss Meditec, Opus Genetics, Viatris, Allergan, Astellas, Alcon, Radius XR, iCare, Glaukos, Eyenovia, Tarsus, Orasis, Topcon, Envision Health Technologies, LKC
- Shareholder-Clearside Biomedical, Annexon Bio (<0.01% ownership)
- *All relevant relationships have been mitigated*

DISCLOSURES FOR DR SCHMIDT


- Dr Schmidt is a consultant or advisor for the following:
 - Tarsus
 - Allergan
 - B&L
 - Tenpoint Pharmaceuticals
 - Topcon
 - Orasis
 - Glaukos
 - Sun Pharmaceuticals
 - Thea Pharmaceuticals
 - Sight Science
 - Alcon
 - Apellis
 - Sydnexis
 - Visus
 - Harrow Pharmaceuticals
 - Lenz Therapeutics
- All relevant financial relationships have been mitigated

64 year old female

Diagnosed with POAG in her 30s with disease progression until IOP within 7-9mmHg range

Netarsudil-latanoprost QHS OU, dorzolamide-timolol BID OU

CCT 477 μ m OD 495 μ m OS





Fixation Monitor: Blind Spot
 Fixation Target: Central
 Fixation Losses: 8/12 XX
 False POS Errors: 9%
 False NEG Errors: 17%
 Test Duration: 03:33
 Fovea: 35 dB

Stimulus: III, White
 Background: 31.5 asb
 Strategy: SITA Faster
 Pupil Diameter:
 Visual Acuity:
 Rx: +3.00 DS

Date: Jan 26, 2022
 Time: 10:37 AM
 Age: 60

*** Low Test Reliability ***



Total Deviation

Pattern Deviation

GHT: Outside Normal Limits



VFI24-2C: 86%
 MD24-2C: -4.91 dB P < 0.5%
 PSD24-2C: 5.67 dB P < 0.5%

:: P < 5%
 :: P < 2%
 :: P < 1%
 ■ P < 0.5%

Fixation Monitor: Gaze Monitor
 Fixation Target: Central
 Fixation Losses: 0/0
 False POS Errors: 9%
 False NEG Errors: 14%
 Test Duration: 03:04
 Fovea: 35 dB

Stimulus: III, White
 Background: 31.5 asb
 Strategy: SITA Faster
 Pupil Diameter: 4.7 mm *
 Visual Acuity:
 Rx: +3.00 DS

Date: Jan 26, 2022
 Time: 10:31 AM
 Age: 60



Total Deviation

Pattern Deviation

GHT: Outside Normal Limits

VFI24-2C: 88%
 MD24-2C: -4.60 dB P < 1%
 PSD24-2C: 4.97 dB P < 0.5%

:: P < 5%
 :: P < 2%
 :: P < 1%
 ■ P < 0.5%

64 year old female

Diagnosed with POAG in her 30s with disease progression until IOP within 7-9mmHg range

Netarsudil-latanoprost QHS OU, dorzolamide-timolol BID OU

CCT 477 μ m OD 495 μ m OS

Is Corneal Thickness an Independent Risk Factor for Glaucoma?

Felipe A. Medeiros, M.D., Ph.D. and Robert N. Weinreb, M.D.

Hamilton Glaucoma Center, University of California San Diego, La Jolla, CA


Ophthalmology. 2012 March ; 119(3): . doi:10.1016/j.ophtha.2012.01.018.

In conclusion, the results of Brandt et al suggest that the use of CCT correction formulas for GAT measurements is probably of little value in clinical practice. Instead of attempting to use these formulas, clinicians are probably better off incorporating risk information as provided by validated predictive models for glaucoma development.^{6, 7} However, the conclusion that CCT is a true independent risk factor for glaucoma is not validated at this time and requires further investigations.

Intraocular Pressure, Central Corneal Thickness, and Prevalence of Open-Angle Glaucoma: The Los Angeles Latino Eye Study

Brian A. Francis, MD, MS¹, Rohit Varma, MD, MPH^{1,2}, Vikas Chopra, MD¹, Mei-Ying Lai, MS², Corina Shtir, MS², Stanley P. Azen, PhD², and Los Angeles Latino Eye Study Group³

Am J Ophthalmol. 2008 November ; 146(5): 741–746. doi:10.1016/j.ajo.2008.05.048.



Decoding the Cornea-Glaucoma Association: Evidence From Mendelian Randomization

Victor A. de Vries,^{1,2} Anita Szabo,³ Joëlle E. Vergroesen,^{1,2} Siyin Liu,^{3,4}
Karin A. van Garderen,^{1,2} Kirithika Muthusamy,⁴ Petra Liskova,⁵ Lubica Dudakova,⁵
Anthony P. Khawaja,⁶ Stephen J. Tuft,^{3,4} Caroline C. W. Klaver,^{1,2,7,8} Alice E. Davidson,^{3,4} and
Wishal D. Ramdas¹

CONCLUSIONS. We found no evidence for a causal link between CCT and OAG. Nevertheless, CCT measurements are still valuable for population-based risk stratification. We found no clear relationship between FECD and OAG.

64 year old female

Recently underwent lower lid blepharoplasty

Reports ongoing left ocular irritation > 6 weeks post-operatively

Surgeon suggested discontinuing dorzolamide-timolol


64 year old female

***IOP 11 mmHg OD and OS (netarsudil-
latanoprost QHS OU)***

Advocate for SLT...anticipated effect?

iCare Home2

Measurement history



64 year old female

Now what?!

***Restart dorzolamide-timolol BID OU; SLT
OU***


Looking ahead...


WHAT THE FACE?!?!?

- 46 y/o WF
- No complaints
- Post Lasik x 12 yrs- stable RE
- Referred because of suspicious disks (by a new eye doctor)

EXAM SPECIFICS

- VA – 20/20 OD, OS w/out RX
- Slit lamp exam – Well apposed LASIK flaps OU
- Gonio – Gr 4 360 degrees OU
- IOP – 14mm Hg OD, 18mm Hg OS
- Pachymetry - 456 OD, 442OS
- Normal Fundus Exam – Except for...










SO WHAT NOW DR GENIUS??

- Any Questions?
- Obviously more testing is needed
- But WTF??



ONH and RNFL OU Analysis: Optic Disc Cube 200x200 OD ● ● OS




	OD	OS
Average RNFL Thickness	98 μm	87 μm
RNFL Symmetry	85%	
Rim Area	1.18 mm^2	0.72 mm^2
Disc Area	2.22 mm^2	2.26 mm^2
Average C/D Ratio	0.70	0.83
Vertical C/D Ratio	0.73	0.86
Cup Volume	0.606 mm^3	0.956 mm^3




Disc Center(-0.15,0.12)mm
Extracted Horizontal Tomogram






Disc Center(0.09,0.00)mm
Extracted Horizontal Tomogram




Extracted Vertical Tomogram




RNFL Circular Tomogram



Extracted Vertical Tomogram




RNFL Circular Tomogram



Fixation Monitor: Gaze Monitor
 Fixation Target: Central
 Fixation Losses: 0/0
 False POS Errors: 0%
 False NEG Errors: 0%
 Test Duration: 02:07
 Fovea: Off


Stimulus: III, White
 Background: 31.5 asb
 Strategy: SITA Faster
 Pupil Diameter: 5.3 mm *
 Visual Acuity:
 Rx: +1.75 DS

Date: Dec 12, 2022
 Time: 4:27 PM
 Age: 52




0 0 0 3
 -2 -1 0 1 2 2
 -2 -3 -2 -2 0 2 -1 1
 -1 -2 -2 -1 -1 1 1 0
 -2 -3 -3 -2 -1 -1 -2 -2
 -2 -6 -3 -1 -2 -4 -1 1
 2 -3 -1 -1 -1 2
 -4 0 0 0

Total Deviation



-1 -1 -1 2
 -3 -2 -2 0 0 0
 -3 -5 -3 -4 -1 0 -2 0
 -3 -3 -3 -2 -2 -1 0 -1
 -3 -4 -4 -3 -2 -2 -3 -3
 -4 -7 -4 -3 -3 -5 -2 0
 1 -5 -3 -2 -2 0
 -5 -1 -1 -1

Pattern Deviation



GHT: **Borderline**

VFI: 98%
 MD24-2: -1.10 dB
 PSD24-2: 1.79 dB P < 10%

:: P < 5%
 P < 2%
 P < 1%
 P < 0.5%

Fixation Monitor: Gaze Monitor
 Fixation Target: Central
 Fixation Losses: 0/0
 False POS Errors: 0%
 False NEG Errors: 0%
 Test Duration: 02:03
 Fovea: Off


Stimulus: Ill, White
 Background: 31.5 asb
 Strategy: SITA Faster
 Pupil Diameter: 4.8 mm *
 Visual Acuity:
 Rx: +1.75 DS

Date: Dec 12, 2022
 Time: 4:24 PM
 Age: 52




1	-3	1	-4
-5	-2	-2	-1
2	-4	-2	-4
0	-3	-2	-4
0	-5	-2	-1
-1	-1	-4	-2
-2	-2	-1	-3
-1	0	0	1

Total Deviation



1	-2	1	-4
-5	-1	-1	-1
3	-4	-2	-3
0	-3	-2	-4
0	-5	-2	-1
0	-1	-4	-2
-1	-1	-1	-3
-1	0	0	1

Pattern Deviation



GHT: Outside Normal Limits

VFI: 97%

MD24-2: 2.26 dB P < 5%

PSD24-2: 2.20 dB P < 5%

:: P < 5%
 :: P < 2%
 :: P < 1%
 ■ P < 0.5%



SO GIVEN THE EVIDENCE BEFORE YOU WHAT DO YOU DO?

- Observe
 - Additional Testing
 - Treat
 - If so, with what?
-
- This is a difficult case, am I right?

WHAT ARE THE BIGGEST “PERTINENTS” IN THIS CASE?

- Optic Nerves?
- OCT?
- Pachymetry?


IF YOU DECIDE TO TREAT...

- What is your Target IOP?
- How Low Do You Go? (How do you know?)
- How do you get the IOP to that Target?





ID: CZMI247065527 Exam Date: 03/13/2023 03/13/2023 CZM
DOB: 08/11/1970 Exam Time: 3:38 PM 3:34 PM
Gender: Female Serial Number: 4000-7182 4000-7182
Technician: Operator, Cirrus Signal Strength: 6/10 7/10

ONH and RNFL OU Analysis: Optic Disc Cube 200x200



OD ● ● OS









	OD	OS
Average RNFL Thickness	88 μm	80 μm
RNFL Symmetry	51%	
Rim Area	0.79 mm ²	0.65 mm ²
Disc Area	2.24 mm ²	2.37 mm ²
Average C/D Ratio	0.81	0.66
Vertical C/D Ratio	0.78	0.66
Cup Volume	0.878 mm ³	1.093 mm ³



Disc Center(-0.12,-0.03)mm
Extracted Horizontal Tomogram




Disc Center(0.21,-0.27)mm
Extracted Horizontal Tomogram



ID: CZMI247065527 Exam Date: 08/08/2019 03/13/2023 CZM
DOB: 08/11/1970 Exam Time: 5:59 AM 3:38 PM
Gender: Female Serial Number: 4000-7182 4000-7182
Technician: Operator, Cirrus Signal Strength: 7/10 6/10

Guided Progression Analysis: (GPA™)

OD ☒ OS ☐



Name: Kern, Maryn

ID: CZMI247065527

DOB: 08/11/1970

Gender: Female


Technician: Operator, Cirrus

Exam Date: 08/08/2019 03/13/2023 CZM

Exam Time: 5:59 AM 3:38 PM

Serial Number: 4000-7182 4000-7182

Signal Strength: 7/10 6/10

**Guided Progression Analysis: (GPA™)**OD ☒ OS ☐

RNFL and ONH Summary Parameters

		Exam Date/Time	Serial Number	Registration Method	SS	Avg RNFL Thickness (µm)	Inf Quadrant RNFL (µm)	Sup Quadrant RNFL (µm)	Rim Area (mm²)	Average Cup-to-Disc Ratio	Vertical Cup-to-Disc Ratio	Cup Volume (mm³)
Baseline1:	1	08/08/2019 5:59:13 AM	4000-7182		7/10	98	120	125	1.18	0.70	0.73	0.606
Baseline2:	2	08/17/2020 12:04:18 PM	4000-7182	R1	7/10	91	111	107	0.72	0.83	0.80	0.931
Current:	3	03/13/2023 3:38:20 PM	4000-7182	R2	6/10	88	108	101	0.79	0.81	0.78	0.878

Registration Methods

R2 - Registration based on translation and rotation of OCT fundus

R1 - Registration based only on translation of disc center

ID: CZMI247065527

DOB: 08/11/1970

Gender: Female

Technician: Operator, Cirrus

Exam Date: 08/08/2019

Exam Time: 6:27 AM

Serial Number: 4000-7182

Signal Strength: 7/10

03/13/2023

3:34 PM

4000-7182





7/10

CZM

Guided Progression Analysis: (GPA™)

OD ☐


OS ☒



ID: CZMI247065527 Exam Date: 08/08/2019 03/13/2023 CZM
DOB: 08/11/1970 Exam Time: 6:27 AM 3:34 PM
Gender: Female Serial Number: 4000-7182 4000-7182
Technician: Operator, Cirrus Signal Strength: 7/10 7/10

Guided Progression Analysis: (GPA™)

OD ☐ OS ☒



RNFL and ONH Summary Parameters

		Exam Date/Time	Serial Number	Registration Method	SS	Avg RNFL Thickness (µm)	Inf Quadrant RNFL (µm)	Sup Quadrant RNFL (µm)	Rim Area (mm²)	Average Cup-to-Disc Ratio	Vertical Cup-to-Disc Ratio	Cup Volume (mm³)
Baseline1:	1	08/08/2019 6:27:51 AM	4000-7182		7/10	87	99	116	0.72	0.83	0.86	0.956
Baseline2:	2	08/17/2020 12:05:44 PM	4000-7182	R2	7/10	85	92	110	0.71	0.83	0.86	1.000
	3	04/12/2021 3:57:44 PM	4000-7182	R2	6/10	81	83	106	0.65	0.86	0.86	1.074
Current:	4	03/13/2023 3:34:32 PM	4000-7182	R2	7/10	80	79	105	0.65	0.86	0.86	1.093

Registration Methods

R2 - Registration based on translation and rotation of OCT fundus
R1 - Registration based only on translation of disc center

Likely Loss

Possible Loss

Compared to baseline, statistically significant loss of tissue detected. For Average RNFL, Superior RNFL, Inferior RNFL, Rim Area the values have decreased. For Cup-to-Disc Ratios and Cup Volume values have increased.

-2.00 D pre-LASIK

Vitreous, alphas p-blurred vision
Azopt - ineffective corneal pt-blurred vision & burning

Notes: LT6 Dx
Rocklatan - Hyperand

GLAUCOMA TRACKING
SHEET

Pachymetry: Date 4/3/17
OD 456 OS 442 - Post Lasik

NAME Marilyn Kern DOB 8/11/70 FHX (+) Father

Date	IOP	Target	C/D	DIL	Gonio	VF	URT	FP	VA	Ref.	Sx	Meds
10/21/16	15.4/18.3	14	151/85 181/8						20/20 20/20	✓	Lasik 1999	Lumi. QHS OU
1/16/17	13/11		151/85 151/9			✓			20/20 20/25			Cont. Lumigan QHS OU
4/3/17	12/12		151/85 151/2				✓		20/20-2 20/20-			Latanoprost
7/11/17	13.8/15.9	12	151/85 151/9	✓				✓	20/20 20/20			Cont. Latanoprost QHS OU
10/12/17	13.7/15	12							20/25+2 20/25-2	✓		D/C Latan. Start Zioptan. QHS OU
1/23/18	16.6/18.7	12	191/88 191/85			✓			20/20- 20/25			D/C Zioptan. Start Lumigan OU
3-26-18	15.1/15.7		191/85 191/9	✓					20/20 20/20-2			Cont. Lumigan QHS OU
6-28-18	12/12						✓		20/20- 20/20-			cont. Lumigan QHS OU
9-24-18	10/11		191/9 191/9						20/20 20/30+	✓		"
1-3-18	10/11		191/9 191/9			✓						Lumigan QHS
5/2/19	12/13		191/90 191/95	✓					N/A			"
8/8/19	12/13	12	191/90 191/90				✓		20/20 20/25-			"
11/7/19	14/13	12	191/85 191/9						20/20 20/20	✓		"
02/10/20	12/12	10	191/85 191/9			✓			20/20 20/20			Rocklatan QHS
06/17/20	11/11	(N)				✓			20/20 20/25			duRocklatan QHS
8/17/20	14/13	10	191/9 191/9			✓			20/20 20/20			d/c Azopt add Alphaganp OU BID OU BID, cont Lumigan QHS
11/9/20	12/11	12	191/9 191/9			✓			20/20 20/20			Cont Lumigan OU QHS
2/8/21	11/12								20/20 20/20			Cont Lumigan OU QHS
4/12/21	12/12	12	191/9 191/9				✓		20/20 20/20			Lumigan OU QHS
7/19/21	12/14		191/85 191/85						20/20 20/25	✓		Lumigan OU QHS
10/21/21	14/13	12	191/85 191/85	✓					20/20 20/21			"Lumigan" OU QHS
1/31/22	12/13	12	191/85 191/85			✓			20/20 20/25			Cont Lumigan OU QHS
8-18-22	13/12	12	191/85 191/85					✓	20/20 20/20			Cont. Lumigan OU QHS
9/26/22	14/14	12	191/85 191/85							✓		Cont. Lumigan
3/13/23	7/17	12	191/9			✓			20/20 20/20			Cont. Lumigan

full

on Virealta

Rocklatan QHS
duRocklatan QHS
d/c Azopt
add Alphaganp OU BID
OU BID, cont Lumigan QHS
Cont Lumigan OU QHS
Cont Lumigan OU QHS
Lumigan OU QHS
Lumigan OU QHS
"Lumigan" OU QHS
Cont Lumigan OU QHS
Cont. Lumigan OU QHS
Cont. Lumigan
okay to use Lumigan
OU BID

IS SHE REALLY STABLE?

- Look at the VF
- Look at the OCT Progression Analysis
- Look at the IOP variability
- How can you maintain a stable and (very) low IOP?

57 year old female



***Peak untreated IOP 21mmHg OD 26mmHg
OS***

***Transferred care for 5 years; returned taking
latanoprost QHS OU...misses about 2 night
per week due to irritation***

“Do I have glaucoma?”

“Do I need to keep taking these eyedrops?”

Adherence during known monitoring



Sustained delivery options

Bimatoprost implant 10mcg Equivalent to about 2-3 drops of bimatoprost 0.01%
Drug release complete in 3-4 months



197 eyes, 94.9% pseudophakic, 41.6% prior SLT, mean age 80.4
Effect approximately 1 year; reduction in medication use
16.9% underwent SLT within first 12 months

Travoprost titanium implant

FDA approved December 14, 2023
Not refillable

36 month data: 70% (fast-release) and **68% (slow-release)** fewer or same medications as baseline

Mean IOP reduction: 8.3mmHg (fast-release) and 8.5mmHg (slow-release)



“Do I really need to use these eyedrops?”

***In office: 16mmHg OD and 17mmHg OS
CCT 555 μ m OD 562 μ m OS***

Medication holiday

IOP 21mmHg OD and 22mmHg OS

OS Single Field Analysis

Central 24-2C Threshold Test



OD Single Field Analysis

Central 24-2 Threshold Test

Fixation Monitor: Gaze Monitor
 Fixation Target: Central
 Fixation Losses: 0/0
 False POS Errors: 0%
 False NEG Errors: 0%
 Test Duration: 02:20
 Fovea: 35 dB

Stimulus: III, White
 Background: 31.5 asb
 Strategy: SITA Faster
 Pupil Diameter: 6.5 mm *
 Visual Acuity:
 Rx: +2.00 DS

Date: Jan 29, 2026
 Time: 11:29 AM
 Age: 57



GHT: Within Normal Limits



VFI24-2C: 99%
 MD24-2C: -1.00 dB
 PSD24-2C: 1.34 dB

:: P < 5%
 :: P < 2%
 :: P < 1%
 ■ P < 0.5%

Fixation Monitor: Gaze Monitor
 Fixation Target: Central
 Fixation Losses: 0/0
 False POS Errors: 0%
 False NEG Errors: 0%
 Test Duration: 02:00
 Fovea: 34 dB

Stimulus: III, White
 Background: 31.5 asb
 Strategy: SITA Faster
 Pupil Diameter: 6.6 mm *
 Visual Acuity:
 Rx: +1.50 DS








Date: Jan 29, 2026
 Time: 11:25 AM
 Age: 57




GHT: Within Normal Limits


VFI24-2: 99%
 MD24-2: -0.24 dB
 PSD24-2: 1.35 dB



:: P < 5%
 :: P < 2%
 :: P < 1%
 ■ P < 0.5%





2019





57 year old female

Primary open angle glaucoma, left eye

Treatment options?

Primary SLT? Latanoprost 0.005% (BAK-preserved, preservative-free)



AMERICAN ACADEMY
OF OPHTHALMOLOGY®



Efficacy of Repeat Selective Laser Trabeculoplasty in Medication-Naive Open-Angle Glaucoma and Ocular Hypertension during the LiGHT Trial

Selective laser trabeculoplasty versus eye drops for first-line treatment of ocular hypertension and glaucoma (LiGHT): a multicentre randomised controlled trial



Gus Gazzard, Evgenia Konstantakopoulou, David Garway-Heath, Anurag Garg, Victoria Vickerstaff, Rachael Hunter, Gareth Ambler, Catey Bunce, Richard Wormald, Neil Nathwani, Keith Barton, Gary Rubin, Marta Buszewicz, on behalf of the LiGHT Trial Study Group*



No game-changing data; But did provide good quality evidence for what was already known

**SLT: at target 93% of visits, 91.3% in the medication group (3 years)
What happens if one eye is drop-free, but the other isn't?**



Laser in Glaucoma and Ocular Hypertension (LiGHT) Trial

Six-Year Results of Primary Selective Laser Trabeculoplasty versus Eye Drops for the Treatment of Glaucoma and Ocular Hypertension

Conclusions: Selective laser trabeculoplasty is a safe treatment for OAG and OHT, providing better long-term disease control than initial drop therapy, with reduced need for incisional glaucoma and cataract surgery over 6 years. *Ophthalmology* 2023;130:139-151 © 2022 by the American Academy of Ophthalmology. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

No difference in health-related quality of life scores: mobility, self-care, usual activities, pain or discomfort.

Minimal difference in Glaucoma Symptom Scale-did not translate to difference in HRQoL



AMERICAN ACADEMY
OF OPHTHALMOLOGY®

Six-Year Rate of Visual Field Progression in the Laser in Glaucoma and Ocular Hypertension Trial

Giovanni Montesano, MD, PhD,^{1,2} David P. Crabb, PhD,² David F. Garway-Heath, MD, FRCOphth,¹ David M. Wright, PhD,³ Evgenia Konstantakopoulou, PhD,^{1,4} Neil Nathwani, BSc (Hons), DipTp(IP),¹ Giovanni Ometto, PhD,^{1,2} Gus Gazzard, MD, FRCOphth¹

Results: Data from 710 eyes (482 with OAG and 354 in the SLT-first arm) were analyzed. The 2 arms had similar baseline MD ($P = 0.7$). The average intraocular pressure (IOP) during follow-up was 16.1 [14.2–18.2] for the drops-first arm and 16.8 [14.6–18.6] in the SLT-first arm (median [interquartile range], $P = 0.057$). The mean [95% credible interval] MD rate was -0.37 [-0.43 to -0.31] decibels (dB)/year in the drops-first arm and -0.26 [-0.31 to -0.21] dB/year in the SLT-first arm ($P = 0.007$). When stratified by severity, this difference was significant only in mild OAG ($P = 0.035$, the largest sub-group). The secondary analyses largely confirmed the main results. The difference in MPD rate was also significantly slower in the SLT-first arm ($P < 0.001$).

26% of med-first eyes were fast-progressors vs. 15% in SLT-first

Selective Laser Trabeculoplasty After Medical Treatment for Glaucoma or Ocular Hypertension

Evgenia Konstantakopoulou, PhD; Gus Gazzard, MA, MD, MBBChir; David Garway-Heath, MD; Mariam Adeleke, PhD; Gareth Ambler, PhD; Victoria Vickerstaff, PhD; Catey Bunce, DSc; Neil Nathwani, BSc; Keith Barton, MD; for the LiGHT Trial Study Group

Patients were randomized to SLT or medication first (LiGHT)-then after 3 years patients were allowed to have secondary SLT (to reduce medication load) or to escalate therapy if needed

65% of 320 taking drops chose to continue medical therapy!

Patients “trust” eyedrops and prefer non-procedures—even though they don’t take them as much as we would like...

57 year old female

Opts for preservative-free latanoprost 0.005%

Case 2 - A Tough Inheritance

- 62 y/o BF, (+) fam hx- treated for POAG for 6 years
- VA 20/20 OD, 20/20 OS
- Pachs – OD 490, OS 495
- No systemic meds
- IOP maintained around 18 OU on Lumigan QHS, AlphaganP OU TID, T1/2 OU BID
- Initial IOP 28 OD, 29OS
- Condition was stable but px developed hypersensitivity (After patient was switched to Brimonidine 0.15%)
- IOP 22 OU on Lumigan only

0000



6000



ID: 100105

CENTRAL 24-2 THRESHOLD TEST



FIXATION MONITOR: GAZE/BLIND SPOT
 FIXATION TARGET: CENTRAL
 FIXATION LOSSES: 1/15
 FALSE POS ERRORS: 2 %
 FALSE NEG ERRORS: 1 %
 TEST DURATION: 05:37

FOVEA: OFF


STIMULUS: III, WHITE
 BACKGROUND: 31.5 ASB
 STRATEGY: SITA-STANDARD

PUPIL DIAMETER:
 VISUAL ACUITY:
 RX: +4.75 DS DC X

DATE: 11-08-2016
 TIME: 9:07 AM
 AGE: 75



TOTAL DEVIATION



PATTERN DEVIATION

GHT

OUTSIDE NORMAL LIMITS

VFI 95%

MD -2.85 DB P < 2%

PSD 3.00 DB P < 2%

11 < 5%
 12 < 2%
 13 < 1%
 14 < 0.5%

BLADEN EYE CENTER
 ERIC SCHMIDT, O.D., /M. SUTTON SCHILAWSKI
 489 E. BROAD ST
 ELIZABETHTOWN NC 28337
 910.862.4268

ID: 100105

CENTRAL 24-2 THRESHOLD TEST



FIXATION MONITOR: GAZE/BLIND SPOT
 FIXATION TARGET: CENTRAL
 FIXATION LOSSES: 4/16 XX
 FALSE POS ERRORS: 1 %
 FALSE NEG ERRORS: 4 %
 TEST DURATION: 07:23

FOVEA: OFF


STIMULUS: III, WHITE
 BACKGROUND: 31.5 ASB
 STRATEGY: SITA-STANDARD

PUPIL DIAMETER:
 VISUAL ACUITY:
 RX: +4.00 DS DC X

DATE: 11-08-2016
 TIME: 8:54 AM
 AGE: 75



TOTAL DEVIATION



PATTERN DEVIATION

*** LOW TEST RELIABILITY ***

GHT

OUTSIDE NORMAL LIMITS

VFI 94%

MD -6.69 DB P < 0.5%

PSD 7.12 DB P < 0.5%

11 < 5%
 12 < 2%
 13 < 1%
 14 < 0.5%

BLADEN EYE CENTER
 ERIC SCHMIDT, O.D., /M. SUTTON SCHILAWSKI
 489 E. BROAD ST
 ELIZABETHTOWN NC 28337
 910.862.4268


Right / OD

Nerve Fiber ONH/GCC OU Report

Left / OS


Exam Date:2015-12-08 09:49:33

Optic Nerve Head Map




Good 64 Scan Quality Index Good 60

RNFL Analysis (μm)		OD	OS	Inter Eye Diff
Avg RNFL Thick...	45	89	-44	
Avg Superior R...	45	88	-43	
Avg Inferior RNFL	45	90	-45	
Intra Eye Diff (S...	0	-2	N/A	
ONH Analysis				
	OD	OS	Inter Eye Diff	
Area C/D	0.24	0.06	0.18	
V. C/D	0.46	0.50	-0.04	
H. C/D	0.53	0.12	0.41	
Rim Area (mm²)	2.97	3.15	-0.18	
Disc Area (mm²)	3.90	3.36	0.54	
Cup Vol (mm³)	0.012	0.000	0.012	



NDB Reference





Exam Date:2015-12-08 09:47:18

Good 76 Scan Quality Index Good 81

GCC Avg Thickness (μm)			
OD	OS	Inter Eye Diff	
Total	69 52	17	
Superior	72 44	28	
Inferior	65 61	4	
Intra Eye Difference (...)	7 -17	N/A	
FLV (%)	6.651 33.2...	-26.596	
GLV (%)	25.6... 44.9...	-19.291	

Exam Date:2015-12-08 09:52:58

Optic Nerve Head Map



NDB Reference



Exam Date:2015-12-08 09:52:07

Report Date: Tuesday 09/24/2019 13:58:40


Software Version: 3.2.1.8

Comment:

Signature:


Attendance Question

- What is the target IOP?
 - 1. ~18
 - 2. ~15
 - 3. ~12




What would you recommend?

- 1. Switch to Rocklatan
- 2. SLT OU 180
- 3. Add Azopt OU BID
- 4. add Timoptic ½ OU BID
- 5. Trabeculectomy
- 6. d/c Lumigan, try Travatan Z OU QHS
- 7. Cosopt OU BID
- 8. Combigan OU BID




SLT OU IOP 19 OD, 20 OS

- What would you do now?



How Do You Know if the IOP needs to be lower?

- What are the risk factors for progression?
 - Age
 - IOP at diagnosis
 - Neuroretinal rim tissue
 - Disk hemes
 - Corneal hysteresis
- Is she progressing?




Right / OD

Nerve Fiber ONH/GCC OU Report

Left / OS

Exam Date:2019-01-30 08:19:20

Optic Nerve Head Map




Good 51


Scan Quality Index

Good 54

RNFL Analysis (μm)		OD	OS	Inter Eye Diff
Avg RNFL Thick...		74	70	4
Avg Superior R...		82	77	5
Avg Inferior RNFL		66	62	4
Intra Eye Diff (S...		16	15	N/A
ONH Analysis		OD	OS	Inter Eye Diff
Area C/D		0.74	0.53	0.21
V. C/D		0.89	0.76	0.13
H. C/D		0.90	0.70	0.20
Rim Area (mm²)		1.03	1.56	-0.53
Disc Area (mm²)		3.94	3.31	0.63
Cup Vol (mm³)		1.348	0.229	1.119



NDB Reference



Exam Date:2019-01-30 08:18:19

Good 65



Scan Quality Index

Good 76


GCC Avg Thickness (μm)		OD	OS	Inter Eye Diff
Total		74	75	-1
Superior		78	77	1
Inferior		70	73	-3
Intra Eye Difference (...)		8	4	N/A
FLV (%)		6.150	11.2...	-5.115
GLV (%)		19.9...	18.6...	1.281

Exam Date:2019-01-30 08:20:53

Optic Nerve Head Map







NDB Reference



Exam Date:2019-01-30 08:20:01





Nerve Fiber ONH/GCC Change Analysis

Left / OS



Nerve Fiber ONH/GCC Change Analysis

Right / OD



NAME: MCALLISTER MATTIE
ID: 100105
DOB: 03-02-1941

CENTRAL 24-2 THRESHOLD TEST



FIXATION MONITOR: BLIND SPOT
FIXATION TARGET: CENTRAL
FIXATION LOSSES: 0/10
FALSE POS ERRORS: 0 %
FALSE NEG ERRORS: 0 %
TEST DURATION: 03:32

FOVER: OFF


STIMULUS: III, WHITE
BACKGROUND: 31.5 ASB
STRATEGY: SITA-FAST

PUPIL DIAMETER:
VISUAL ACUITY:
RX: +4.00 DS DC X


DATE: 04-02-2019
TIME: 9:27 AM
AGE: 70



TOTAL DEVIATION



PATTERN DEVIATION



CWT
OUTSIDE NORMAL LIMITS

VF1 94%
MD -2.38 DB P < 5%
PSD 3.71 DB P < 0.5%

BLADE EYE CENTER
ERIC SCHMIDT, O.D., F.N. SUTTON SCHILAWSKI
409 E. BROAD ST
ELIZABETHTOWN NC 28337
910.662.4268

NAME: MCALLISTER MATTIE
ID: 100105

CENTRAL 24-2 THRESHOLD TEST



FIXATION MONITOR: GAZE/BLIND SPOT
FIXATION TARGET: CENTRAL
FIXATION LOSSES: 4/12 XX
FALSE POS ERRORS: 0 %
FALSE NEG ERRORS: 2 %
TEST DURATION: 04:01

FOVER: OFF


STIMULUS: III, WHITE
BACKGROUND: 31.5 ASB
STRATEGY: SITA-FAST

PUPIL DIAMETER:
VISUAL ACUITY:
RX: +3.75 DS DC X


DATE: 04-02-2019
TIME: 9:21 AM
AGE: 70



TOTAL DEVIATION



PATTERN DEVIATION



*** LOW TEST RELIABILITY ***

CWT
OUTSIDE NORMAL LIMITS

VF1 83%
MD -6.29 DB P < 0.5%
PSD 7.33 DB P < 0.5%

BLADE EYE CENTER
ERIC SCHMIDT, O.D., F.N. SUTTON SCHILAWSKI
409 E. BROAD ST
ELIZABETHTOWN NC 28337
910.662.4268

So Let's Talk About Compliance



Adherence to IOP-Lowering Therapy


A complex and multifaceted problem^{1,2}

- Includes both persistency and compliance issues

Factors related to adherence¹⁻³

- Problems tolerating therapy
- Safety issues
- Inadequate efficacy
- Convenience factors, eg, access, cost, dosing
- Patient/situational factors, eg:
 - Competing activities
 - Forgetfulness
 - Poor disease knowledge and health literacy

Over 3 months in a study of 196 patients with glaucoma taking an IOP-lowering medication in one or both eyes^{2,4}:



Despite instruction, free medication, once-daily administration, use of a dosing aid, and electronic monitoring of adherence

IOP=intraocular pressure.

Predictors of Poor Adherence – Friedman 2019

- Gaps In Visits
- Patients Don't Understand Severity Of Disease
- Cost of Drops (25%)
- Those who Travel A Lot
- Younger Pxs and Very Old Pxs
- African-Americans
- Those In Poor Health
 - These drop adherence to <60%

Ways To Improve Compliance

- See Pxs more frequently... especially early in treatment
 - Improve tracking system – better identify no shows
 - Call/email appointment reminders
 - Reminders to pxs to take their drops
 - Change Dr/Patient intervention
-
- G pxs ask 3.2 questions at visit whereas in other chronic diseases pxs ask ~ 6 questions/visit

Choosing Therapy: Switch or Add?

- Target IOP should be reached with minimal number of medications
- Build strongest foundation prior to adjunctive therapy
- Switching to a more effective medication may be best course of action
- Differences within medication classes exist
- Potential benefits of monotherapy
 - Improved patient compliance
 - Decreased cost
 - Reduced cumulative exposure to benzalkonium chloride

Estimating the Lead Time Gained by Optical Coherence Tomography in Detecting Glaucoma before Development of Visual Field Defects

Tammy M. Kuang, MD,^{1,2,3} Chunwei Zhang, MD,^{1,4} Linda M. Zangwill, PhD,¹ Robert N. Weinreb, MD,¹ Felipe A. Medeiros, MD, PhD¹

- At 95% specificity, up to **35% of eyes had abnormal average RNFL thickness** 4 years before development of visual field loss and **19% of eyes had abnormal results 8 years before field loss**.
- **Conclusions:** Assessment of RNFL thickness with OCT was able to detect glaucomatous damage before the appearance of VF defects on SAP. In many subjects, significantly large lead times were seen when applying OCT as an ancillary diagnostic tool.

So using an OCT;


How do we tell if they are getting worse?

- Progression Analysis Software!!!!!!




Guided Progression Analysis: (GPA™)

OD ☐ OS ☒



A NEW LOOK AT GLAUCOMA PROGRESSION


- WHY DO SOME PATIENTS PROGRESS AND SOME DON'T?
- WHY DO SOME PROGRESS AT A FASTER RATE?



Choosing Therapy: Switch or Add?

- Target IOP should be reached with minimal number of medications
- Build strongest foundation prior to adjunctive therapy
- Switching to a more effective medication may be best course of action
- Differences within medication classes exist
- Potential benefits of monotherapy
 - Improved patient compliance
 - Decreased cost
 - Reduced cumulative exposure to benzalkonium chloride

67 year old female





■ What changed?

- Multiple IVI (approximately 6-9 per year) between 2014 and 2016 for macular edema secondary to CRVO
- Bevacizumab, dexamethasone intravitreal implant 0.7mg

HOW *COULD* IVI AND VITRECTOMY CAUSE ELEVATED IOP?!

- Widening indications for vitrectomy
- Anti-VEGF injections are the cornerstone of medical retinal treatment
- **Long and short term IOP rise possible**
- Development of OAG & progression of OAG



SHORT TERM

- We tend to think about the greatest risk of IVI to be endophthalmitis (1/2659)
- Immediately after injection: IOP rise to up to **87mmHg**
 - Most patients increase approximately 20mmHg-35mmHg
 - *Do most surgeons measure IOP after injections?*
- How does this happen?!
 - Increased intravitreal volume
 - 4-4.4mL average volume; most injections 0.05mL; FDA-approved GA therapeutics 0.1mL


Transient vision and intraocular pressure changes following anti-vascular endothelial growth factor injection

Khaldon Abbas,^{*,†} Amy Basilious,^{*,†} Julie Duncan,^{*} Thomas Sheidow,^{*,†} Philip Hooper,^{*,†} John Gonder^{*,†}

At 1 minute after intravitreal injection (IVI), VA worsened significantly from a baseline of 0.29 ± 0.21 to 0.76 ± 0.65 logMAR ($p < 0.001$), while IOP rose from 14.34 ± 4.39 mm Hg to 54.53 ± 20.21 mm Hg ($p < 0.001$). VA progressively improved over time, with 43% of eyes returning to baseline at 1 minute, 67% at 10 minutes, 83% at 20 minutes, and 89% at 30 minutes. The Spearman correlation coefficient for VA and IOP was statistically significant at 1-minute follow-up after IVI at 0.244 ($p = 0.014$) but not at later follow-ups. Eyes returning to baseline VA at 1 minute exhibited lower IOP than those not at baseline ($p = 0.0004$). Logistic regression revealed no significant predictors of delayed VA recovery.

WELL THAT CAN'T BE GOOD

- Risk of retinal artery occlusion (as high as 1/1389 Gao et al 2019)
- Repeated, sudden, **significant** IOP spike and temporary loss of perfusion



HOW CAN WE FIX THIS?

- **Treatment for elevated IOP vs. IOP spike-prevention**

- Role of pre-procedure IOP lowering medication

- Paracentesis

- 32 gauge needle

- Fluid balance

**Anterior chamber paracentesis
during intravitreal injections in observational
trials: effectiveness and safety and effects**



Sandeep Saxena¹, Timothy Y. Lai², Hideki Koizumi³, Michel E. Farah⁴, Daniela Ferrara⁵, David Pelayes⁶,
Tomohito Sato⁷, Carsten H. Meyer^{8*}, Timothy Murray⁹ and for the International Pharmacokinetic Collaboration

ALL ABOUT OUTFLOW

- Reduced trabecular outflow:
 - ~~■ 1) Direct toxicity of medication~~
 - 2) Inflammation
 - Trabeculitis
 - 3) Aggregation of particles
 - Silicone, protein in the TM
 - 4) Nitric oxide reduction

SILICONE?

- Medical grade silicone oil droplets
 - Barrel of the syringe
 - Hub of the needle
 - Tip of the plunger
 - Stopper of the medication vial
- *Silicone oil has the potential to be pro-inflammatory*




NITRIC OXIDE

- Nitric oxide is involved in the signaling pathway which leads to relaxation of trabecular beams
 - Leads to increased trabecular outflow
 - Latanoprostene bunod
 - Latanoprost acid + butanediol monohydrate
 - NO is a gas, so must be attached to another molecule
- VEGF upregulates nitric oxide synthase = increased nitric oxide
- *Effect of **anti-VEGF** medications?*


SO WHO IS MOST AT RISK?

- Greater number of injections (20+)
- Higher frequency of injections (7/year +)
- Younger patients
- Patients with shorter axial length




VITRECTOMY & TAMPONADE AGENTS

- Long term potential for IOP rise
 - Oxidative stress-fluid/air exchange
- Tamponade agents
 - Sulfur hexafluoride (SF_6)
 - Perfluoropropane (C_3F_8)
 - Silicone oil-greatest risk of IOP elevation-as high as 40%



BOTTOM LINE

- **Monitor intraocular pressure in patients undergoing IVI or who have a history of PPV**



A CLASSIC CASE

- 68 Y/O AFRICAN AMERICAN MALE
- PMH: HBP, HYPERCHOLESTEROLEMIA
- FMH: MOTHER POAG
- INITIAL PRESENTATION:

BCVA: 20/20 OD, OS


IOP: OD 21MM HG, OS 22MM HG

C/D: 0.6/0.6 OD, 0.6/0.6 OS

SO WHAT NOW???

OCT 2017

2017





OCT=optical coherence tomography.





Patient ID: ---
Diagnosis: ---

Exam.: May/1/2014
Comment: ---

OD




Asymmetry OD - OS



OS

Classification OD



Classification OS

Within Normal Limits

Handwritten notes in the bottom right corner.

CASE 1 (CONT)

- FP TAKEN
- VF – WNL
- PACHYMETRY – OD 505, OS 510
- HYSTERESIS- OD 8 OS 9
- TO TREAT OR NOT TO TREAT – THAT IS THE QUESTION!!

ARS QUESTION1

- BASED ON THE EVIDENCE AT HAND, WOULD YOU
 1. INITIATE TREATMENT
 2. SEE PATIENT IN 1 MONTH
 3. SEE PATIENT IN 3 MONTHS
 4. SEE PATIENT IN 6 MONTHS
 5. SEE PATIENT IN 12 MONTHS

2 YEARS LATER – NO TREATMENT

2019 – IOP: 20mm Hg OD, 21mm Hg OS

Followed for 2 years w/out tx

IOP ranged 19-23mm Hg (5 readings)


OPTIC NERVE HEAD PHOTO- 2019

VF – WNL OU

OD




OS





OCT 2017 VS 2019

2017





2019



OCT=optical coherence tomography.


VF RESULTS-2019



DEFINITELY POAG

- WERE THERE ANY CLUES THAT MAY HAVE LED US TO TREAT EARLIER??
- WHAT IS THE SIGNIFICANCE OF THE DISK HEMORRHAGES??
- WHAT IS THE SIGNIFICANCE OF THIN CORNEAS??
- WAS THE OCT REALLY NORMAL?:
- TIME FOR A COFFEE TALK!!!

DID YOU SEE THE DISC HEMORRHAGE?



Detection and Prognostic Significance of Optic Disc Hemorrhages during the Ocular Hypertension Treatment Study

Donald L. Budenz, MD, MPH,¹ Douglas R. Anderson, MD,¹ William J. Feuer, MS,¹ Julia A. Beiser, MS,² Joyce Schiffman, MS,¹ Richard K. Parrish II, MD,¹ Jody R. Piltz-Seymour, MD,³ Mae O. Gordon, PhD,² Michael A. Kass, MD,² Ocular Hypertension Treatment Study Group

Main Outcome Measures: Incidence of optic disc hemorrhages and POAG end points.

Results: Median follow-up was 96.3 months. Stereophotography-confirmed glaucomatous optic disc hemorrhages were detected in 128 eyes of 123 participants before the POAG end point. Twenty-one cases (16%) were detected by both clinical examination and review of photographs, and 107 cases (84%) were detected only by review of photographs ($P < 0.0001$). Baseline factors associated with disc hemorrhages were older age, thinner corneas, larger vertical cup-to-disc ratio, larger pattern standard deviation index on perimetry, family history of glaucoma, and smoking status. The occurrence of a disc hemorrhage increased the risk of developing POAG 6-fold in a univariate analysis ($P < 0.001$; 95% confidence interval, 3.6–10.1) and 3.7-fold in a multivariate analysis



Detection and Prognostic Significance of Optic Disc Hemorrhages during the Ocular Hypertension Treatment Study

Donald L. Budenz, MD, MPH,¹ Douglas R. Anderson, MD,¹ William J. Feuer, MS,¹ Julia A. Beiser, MS,² Joyce Schiffman, MS,¹ Richard K. Parrish II, MD,¹ Jody R. Piltz-Seymour, MD,³ Mae O. Gordon, PhD,² Michael A. Kass, MD,² Ocular Hypertension Treatment Study Group

- Disc hemorrhages detected in 128 eyes of 123 participants
- 21 cases detected by both doctor and photos
- **107 cases (84%) were detected only by a review of photography**



DISK HEMORRHAGES AND RATE OF PROGRESSION (MEDEIROS ET AL)

- Cohort of the DIGS
- Pxs followed for 8 years for VF progression (using the VFI)
- 20% had disk hemorrhage
- Eyes with disk heme had more than double the rate of VF loss
- Eyes w/ more than 1 disk heme showed an even higher rate of VF progression
- Persons with disk heme in general had a more severe glaucoma



SPEAKING OF OPTIC DISK HEMORRHAGES

- BUDENZ ET AL, (OHTS GROUP) – AJO 2/17
- 13 YEAR DATA
- ODH ARE AN INDEPENDENT PREDICTOR FOR POAG
- ODH ARE PREDICTIVE OF PROGRESSION
- PREDICTIVE FACTORS FOR ODH ARE SIMILAR TO THOSE FOR POAG (IN OHT PXS)
 - Thin corneas
 - Thinner rims
 - Higher IOP
 - Older age



THE PACHYMETRY ISSUE

➤ JUICY DATA


- 36% OF PXS W/ IOP > 25.75 AND K THICKNESS < 555 MICRONS DEVELOPED POAG
- 6% OF PXS W/ SAME IOP BUT K THICKNESS > 588 CONVERTED TO POAG

➤ JUICY DATA II

- 15% PXS W/ C/D .3/.3 AND K THICKNESS < 555 MICRONS CONVERTED BUT
- 4% OF PXS W/ SAME DISK PARAMETERS AND K THICKNESS > 588 MICRONS CONVERTED


Figure 1

Risk of Progression as a Function of IOP vs. Central Corneal Thickness



The percentage of participants in the observation group who developed primary open-angle glaucoma grouped by baseline IOP and by CCT measurements.

Risk of Progression as a Function of IOP vs. Central Corneal Thickness



The percentage of participants in the observation group who developed primary open-angle glaucoma grouped by vertical cup:disc ratio and by CCT measurements.⁵

IOP 30mmHg, CCT 600μ

Glaucoma Risk Estimator

Age <input type="text" value="70"/>	RIGHT EYE MEASUREMENTS			LEFT EYE MEASUREMENTS		
	1 st	2 nd	3 rd	1 st	2 nd	3 rd
Untreated Intraocular Pressure (mm Hg)	30	30	30	30	30	30
Central Corneal Thickness (microns)	600	600	600	600	600	600
Vertical Cup to Disc Ratio by Contour	0.55			0.55		
Pattern Standard Deviation Humphrey ☒ (dB) Octopus loss variance ☐ (dB)	1.0	1.0		1.0	1.0	

The patient's estimated 5-year risk (%) of developing glaucoma in at least one eye.

Glaucoma risk is 9.1%

IOP 20mmHg, CCT 500μ

Glaucoma Risk Estimator

Age	RIGHT EYE MEASUREMENTS			LEFT EYE MEASUREMENTS		
	1 st	2 nd	3 rd	1 st	2 nd	3 rd
70						
Untreated Intraocular Pressure (mm Hg)	20	20	20	20	20	20
Central Corneal Thickness (microns)	500	500	500	500	500	500
Vertical Cup to Disc Ratio by Contour	0.55			0.55		
Pattern Standard Deviation Humphrey ☒ Octopus ☐ (dB) (dB)	1.0	1.0		1.0	1.0	

[Print](#)[Reset](#)

20.7%

The patient's estimated 5-year risk (%) of developing glaucoma in at least one eye.

Glaucoma risk is 20.7%



So a man walks into his
optometrist's office...

- He is diagnosed with glaucoma,
- What is your initial treatment??

